WEEKLY EPIDEMIOLOGY BULLETIN

NATIONAL EPIDEMIOLOGY UNIT, MINISTRY OF HEALTH, JAMAICA

Weekly Spotlight Stay super!



Quiz: How much do you know about diabetes?

- Obesity increases the risk of getting diabetes.

 - False
- Risk of getting diabetes is higher if a parent has it.
 - True
 - False
- 3 People with type 1 diabetes can live without insulin.
 - True
 - False
- Diabetes can cause kidney failure and blindness.
 - True
 - False
- 5 Regular physical activity has no impact on preventing diabetes or its complications.
 - True
 - False

How well did you do!?



Source: http://who.int/campaigns/world-health-day/2016/quiz/en/



SYNDROMES

WEEK 12

PAGE 2



CLASS 1 DISEASES

PAGE 4



INFLUENZA

PAGE 5



DENGUE FEVER

PAGE 6



GASTROENTERITIS

PAGE 7



NOTIFICATIONS-A11 clinical sites



INVESTIGATION REPORTS- Detailed Follow up for all Class One Events



HOSPITAL ACTIVE SURVEILLANCE-30 sites*. Actively pursued




REPORTS FOR SYNDROMIC SURVEILLANCE


FEVER

Temperature of $>38^{\circ}C$ /100.4°*F* (or recent history of fever) with or without an obvious diagnosis or focus of infection.









FEVER AND NEUROLOGICAL

Temperature of >380C /100.40F (or recent history of fever) in a previously healthy person with or without headache and vomiting. The person must also have meningeal irritation, convulsions, altered consciousness, altered sensory manifestations paralysis (except AFP).





Fever and Neurological Symptoms Weekly Threshold vs Cases 2016, Epidemiology Week 12 30 10 1 3 5 7 9 11 13 15 17 19 21 23 25 27 29 31 33 35 37 39 41 43 45 47 49 51

Epi Weeks

2016

FEVER AND HAEMORRHAGIC


Temperature of $>38^{\circ}C$ /100.4°F (or recent history of fever) in a previously healthy person presenting with at least one haemorrhagic (bleeding) manifestation with or without jaundice.





Fever and Haem Weekly Threshold vs Cases 2016, Epidemiology Week 12

Epidemic Threshold





NOTIFICATIONS-All clinical sites



INVESTIGATION
REPORTS- Detailed Follow
up for all Class One Events



HOSPITAL ACTIVE SURVEILLANCE-30 sites*. Actively pursued




FEVER AND JAUNDICE

Temperature of $>38^{\circ}C$ /100.4°*F* (or recent history of fever) in a previously healthy person presenting with jaundice.








ACCIDENTS

Any injury for which the cause is unintentional, e.g. motor vehicle, falls, burns, etc.








VIOLENCE

Any injury for which the cause is intentional, e.g. gunshot wounds, stab wounds, etc.









NOTIFICATIONS-All clinical sites



INVESTIGATION REPORTS- Detailed Follow up for all Class One Events



HOSPITAL ACTIVE SURVEILLANCE-30 sites*. Actively pursued



CLASS ONE NOTIFIABLE EVENTS

Comments

	CLASS 1 EVENTS		CONFIR	AFP Field Guides		
			CURRENT YEAR	PREVIOUS YEAR	from WHO indicate that for an effective surveillance	
AL	Accidental Poisoning		10	45	system, detection rates for AFP	
NATIONAL /INTERNATIONAL INTEREST	Cholera		0	0	should be	
	Dengue Hemorrhagic Fever ¹		1	0	1/100,000 population under	
L /INTERN INTEREST	Hansen's Disease (Leprosy)		1	0	15 years old (6 to 7) cases annually.	
INTI IER	Hepatitis B		2	14		
	Hepatitis C		0	1		
NO V	HIV/AIDS - See HIV/AIDS National Programme Report				Pertussis-like syndrome and	
ATI	Malaria (Imported)		1	0	Tetanus are	
Z	Meningitis		6	30	clinically confirmed	
EXOTIC/ UNUSUAL	Plague	Plague		0	classifications.	
<u> </u>	Meningococcal Meningitis		0	0	The TB case	
H IGH ORBIDI ORTAL	Neonatal Tetanus		0	0	detection rate	
H IGH MORBIDITA MORTALIY	Typhoid Fever		0	0	established by PAHO for Jamaica	
ΣΣ	Meningitis H/Flu		0	0	is at least 70% of	
	AFP/Polio		0	0	their calculated estimate of cases in	
	Congenital Rubella Syndrome		0	0	the island, this is	
S	Congenital Syphilis		0	0	180 (of 200) cases per year.	
MMES	Fever and	Measles	0	0	per year.	
ZAM	Rash	Rubella	0	0	*Data not available	
[90]	Maternal Deaths ²		14	16		
C PR	Ophthalmia Neonatorum		139	85	1 Dengue Hemorrhagic Fever data include	
CIAI	Pertussis-like syndrome		0	0	Dengue related deaths;	
SPECIAL PROGRA	Rheumatic Fever		0	12	2 Maternal Deaths include early and late	
	Tetanus		0	1	deaths.	
	Tuberculosis		0	0		
	Yellow Fever		0	0		
Chikungunya		ı	0	1		
	Zika Virus		1	0		



All

sites












NATIONAL SURVEILLANCE UNIT INFLUENZA REPORT

March 20- March 26, 2016

Epidemiology Week 12

E	W	12

February, 2016				
	EW 12	YTD		
SARI cases	53	448		
Total Influenza positive Samples	0	97		
Influenza A	0	70		
H3N2	0	1		
H1N1pdm09	0	68		
Not subtyped	0	27		
Influenza B	0	0		
Other	0	1		
a				




Comments:

The percent positivity among all samples tested from EW 1 to EW 8, 2016 is 40.3% (N= 77)

Influenza A(H1N1)pdm09 continued to circulate in EWs 1 to 8 as the predominant virus at 97%. No Influenza B viruses have been detected since 2016. In addition, there has been no detection of the influenza A/H3v or A/H1v variant viruses, or avian H5 and H7 viruses among human samples tested.

Distribution of Influenza and other respiratory viruses by EW surveillance EW 8, 2016, NIC Jamaica - Interim report



INDICATORS

Burden


respiratory date, syndromes account for 5.3% of visits to health facilities.

Incidence

Cannot be calculated, as data sources do not collect all cases of Respiratory illness.

Prevalence

applicable acute respiratory conditions.



*Additional data needed to calculate Epidemic Threshold



NOTIFICATIONS-All clinical sites



INVESTIGATION REPORTS- Detailed Follow up for all Class One Events




HOSPITAL ACTIVE **SURVEILLANCE-30** sites*. Actively pursued




Dengue Bulletin

March 20- March 26, 2016


Epidemiology Week 12



DISTRIBUTION Year-to-Date Suspected Dengue Fever Un-F **Total** M % kwn <1 2 0 1 1-4 0 0 1 0 5-14 2 2 0 4 2 15-24 2 0 3 1 25-44 0 0 1 0 45-64 0 0 0 0 ≥65 0 0 0 0 0 Unknown 138 198 122 458 96 **TOTAL** 143 204 122 469 100



Weekly Breakdown of suspected and confirmed cases of DF,DHF,DSS,DRD 2016 2015 **FW YTD YTD** 12 **Total Suspected** 7 469 24 **Dengue Cases Lab Confirmed** 3 41 1 **Dengue cases DHF/DSS** 0 1 0 CONFIRMED **Dengue** 0 0 Related 0 **Deaths**









INVESTIGATION REPORTS- Detailed Follow up for all Class One Events



HOSPITAL ACTIVE SURVEILLANCE-30 sites*. Actively pursued



Gastroenteritis Bulletin

March 20- March 26, 2016

Epidemiology Week 12

Weekly Breakdown of Gastroenteritis cases

Year	EW 12			YTD		
	<5	≥5	Total	<5	≥5	Total
2016	172	191	363	1941	2724	4665
2015	289	300	589	4259	3884	7843

Figure 1: Total Gastroenteritis Cases Reported 2014-2016


Gastroenteritis: Three or more loose stools within 24 hours.


In Epidemiology Week 12, 2016, the total number of reported GE cases showed a 38% decrease compared to EW 12 of the previous year.

The year to date figure showed a 41% decrease in cases for the period.



















RESEARCH PAPER

A Need for Capacity Building in Faith-Based Response to HIV/AIDS in Jamaica

N Muturi 1, R Page 2

1A.O. Miller School of Journalism and Mass Communications, Kansas State University 2Ministry of Health, Jamaica

Email: nmuturi@ksu.edu or raunchygp@gmail.com

Objective: To identify initiatives being conducted by faith-based organizations (FBOs) and explore their most urgent needs in addressing the HIV/AIDS epidemic.

Design and Methods: Focus group discussions (FGD) and in-depth interviews were conducted with members of FBOs, members of HIV/AIDS support groups and persons living with HIV/AIDS (PLWHA) over a 6 month period in three parishes. Twelve (12) FGD and 30 in-depth interviews were conducted. Data were analysed by descriptive and interpretive techniques following the completion of transcriptions of the interviews and focus groups.

Results: One hundred (100) persons participated in the study, 18 of which were PLWHA. Approximately 60% of FBOs who participated had initiatives to address stigma and discrimination which included education and counselling sessions with their congregants (60%) as well as providing psychological support to PLWHA (50%). One FBO also had media publication. More than 50% of the FBO leaders interviewed expressed their most urgent need to be strengthening of the leadership to address stigma and discrimination and treatment of PLWHA among their congregants.

Conclusions: Programs to address stigma and discrimination were the most common initiatives in the FBOs that participated in the study. Strengthening the capacity of FBO leaders to identify and address stigma and discrimination among their congregants and the wider community was identified as their most urgent need followed by the capacity to provide psychological support for PLWHA.



The Ministry of Health 24-26 Grenada Crescent Kingston 5, Jamaica Tele: (876) 633-7924

Email: mohsurveillance@gmail.com



A11

sites



clinical





